

On the proper choice of a Lorentz-covariant relativistic Lagrangian

Alain J. Brizard

Department of Chemistry and Physics, Saint Michael's College, Colchester, VT 05439, USA

A proper choice for the sign of the Lorentz-covariant relativistic Lagrangian for a particle moving in an electromagnetic field is presented. This choice is based on which space-time metric is used in the representation of the relativistic particle dynamics.

The relativistic Lagrangian for a particle moving in an electromagnetic field is defined up to an overall sign. In almost all textbooks, the Lagrangian is defined with an overall negative sign [1, 2, 3, 4], while in some, the Lagrangian is defined with a positive sign [5, 6]. In the present brief note, we show that the overall sign of the Lorentz-covariant relativistic Lagrangian is determined consistently by the choice of space-time metric used in describing the particle dynamics:

$$\mathbf{g} \equiv \begin{pmatrix} \sigma & 0 & 0 & 0 \\ 0 & -\sigma & 0 & 0 \\ 0 & 0 & -\sigma & 0 \\ 0 & 0 & 0 & -\sigma \end{pmatrix}, \quad (1)$$

where $\sigma = +1$ for a time-like metric and $\sigma = -1$ for a space-like metric.

First, we consider the following relativistic Lagrangian describing the motion of a particle (of mass m and charge e) under the influence of an electromagnetic field

$$L_\epsilon \equiv \epsilon \left[mc^2 \gamma^{-1} + e \left(\Phi - \mathbf{A} \cdot \frac{\dot{\mathbf{x}}}{c} \right) \right], \quad (2)$$

where $\epsilon = \pm 1$ and $A^\mu = (\Phi, \mathbf{A})$ denotes the components of the electromagnetic four-potential. Here, the relativistic factor is

$$\gamma = 1/\sqrt{1 - |\dot{\mathbf{x}}|^2/c^2}, \quad (3)$$

and a dot refers to a time derivative. The choice $\epsilon = -1$ is most commonly adopted because the regular Hamiltonian H is defined by regular Legendre transformation as

$$H \equiv \dot{\mathbf{x}} \cdot \frac{\partial L_-}{\partial \dot{\mathbf{x}}} - L_- = m\gamma c^2 + e\Phi, \quad (4)$$

while the classical Lagrangian

$$L_{\text{cl}} = \left(m\mathbf{v} + \frac{e}{c}\mathbf{A} \right) \cdot \dot{\mathbf{x}} - \left(\frac{m}{2}|\mathbf{v}|^2 + e\Phi \right) \quad (5)$$

is obtained from the relativistic Lagrangian L_- in the nonrelativistic limit ($|\dot{\mathbf{x}}| \ll c$), where the constant $-mc^2$ was omitted from Eq. (5). Here, the identity $\mathbf{v} \equiv \dot{\mathbf{x}}$ for the particle velocity results from the Euler-Lagrange equation for $\mathbf{v}$ (since $\partial L_{\text{cl}}/\partial \dot{\mathbf{v}} \equiv 0$).

Next, we show that the choice for ϵ for a Lorentz-covariant relativistic Lagrangian can be consistently

based on the sign σ of the space-time metric (1). For this purpose, we introduce the notation

$$A^0 B^0 - \mathbf{A} \cdot \mathbf{B} \equiv \sigma A^\mu B_\mu = \sigma A^\mu g_{\mu\nu} B^\nu, \quad (6)$$

where $A^\mu = (A^0, \mathbf{A})$ and $B^\mu = (B^0, \mathbf{B})$ are two arbitrary four-vectors and we used the space-time metric (1). Hence, the relativistic factor (3) is defined covariantly as

$$\gamma \equiv c(\sigma \dot{x}_\mu \dot{x}^\mu)^{-1/2}. \quad (7)$$

The relativistic Lagrangian (2) can be written in Lorentz-covariant form as

$$\begin{aligned} \mathcal{L}_\epsilon &\equiv \epsilon mc \sqrt{(\dot{x}^0)^2 - |\dot{\mathbf{x}}|^2} + \epsilon \frac{e}{c} (\Phi \dot{x}^0 - \mathbf{A} \cdot \dot{\mathbf{x}}) \\ &= \epsilon \left(mc \sqrt{\sigma \dot{x}_\mu \dot{x}^\mu} + \frac{e}{c} \sigma A_\mu \dot{x}^\mu \right) \end{aligned} \quad (8)$$

where $\dot{x}^0 = c$. The action integral

$$S_\epsilon = \int \mathcal{L}_\epsilon dt \equiv \epsilon \int \left(mc ds + \frac{e}{c} \sigma A_\mu dx^\mu \right) \quad (9)$$

therefore involves an integration involving two Lorentz invariants $mc ds$ and $(e/c) A_\mu dx^\mu$, where $ds^2 \equiv \sigma dx_\mu dx^\mu$ and used the definition (6). The Euler-Lagrange equations

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}_\epsilon}{\partial \dot{x}^\mu} \right) = \frac{\partial \mathcal{L}_\epsilon}{\partial x^\mu}$$

obtained from the relativistic Lagrangian (8) or the variational principle based on the action integral (9), are expressed as

$$\frac{d}{dt} \left(mc \gamma \frac{dx_\mu}{dt} + \frac{e}{c} A_\mu \right) = \frac{e}{c} \left(\frac{\partial A_\nu}{\partial x^\mu} \frac{dx^\nu}{dt} \right), \quad (10)$$

where the common factor of $\epsilon \sigma$ has been removed from both sides [i.e., the Euler-Lagrange equations (10) are independent of the choice of the sign ϵ of the Lagrangian and the sign σ of the space-time metric]. If we now define the proper time derivative $d/d\tau \equiv \gamma d/dt$ and the four-velocity $u^\mu \equiv dx^\mu/d\tau$, the Euler-Lagrange equations (10) become (upon multiplication by γ)

$$m \frac{du_\mu}{d\tau} = \frac{e}{c} \left(\frac{\partial A_\nu}{\partial x^\mu} - \frac{\partial A_\mu}{\partial x^\nu} \right) u^\nu \equiv \frac{e}{c} F_{\mu\nu} u^\nu. \quad (11)$$

Note that these equations can be obtained directly from the variational principle $\delta \int \mathcal{L}_\epsilon d\tau = 0$, where

$$\mathcal{L}_\epsilon \equiv \gamma \mathcal{L}_\epsilon = \epsilon \left(mc \sqrt{\sigma u_\mu u^\mu} + \frac{e}{c} \sigma A_\mu u^\mu \right). \quad (12)$$

The associated Euler-Lagrange equations

$$\frac{d}{d\tau} \left(\frac{\partial \mathcal{L}_\epsilon}{\partial u^\mu} \right) = \frac{\partial \mathcal{L}_\epsilon}{\partial x^\mu}$$

directly yield the covariant equations of motion (11).

The covariant and contravariant components of the canonical four-momentum are derived from the Lagrangian (8) as

$$p_0 \equiv \frac{\partial \mathcal{L}_\epsilon}{\partial \dot{x}^0} = \epsilon \left(m \gamma \dot{x}^0 + \frac{e}{c} \Phi \right) \equiv \sigma p^0, \quad (13)$$

$$p_i \equiv \frac{\partial \mathcal{L}_\epsilon}{\partial \dot{x}^i} = -\epsilon \left(m \gamma \dot{x}^i + \frac{e}{c} A^i \right) \equiv -\sigma p^i, \quad (14)$$

where we used the space-time metric (1) to relate the covariant and contravariant components $p_\mu \equiv g_{\mu\nu} p^\nu$. A consistent choice for the sign ϵ of the relativistic Lagrangian (8) [and Eq. (12)] is $\epsilon \equiv \sigma$ and, therefore, the proper choice for the Lorentz-covariant relativistic Lagrangian for a particle moving in an electromagnetic field is

$$\mathcal{L}_\sigma \equiv \sigma mc \sqrt{\sigma \dot{x}_\mu \dot{x}^\mu} + \frac{e}{c} A_\mu \dot{x}^\mu, \quad (15)$$

or

$$\mathcal{L}_\sigma \equiv \sigma mc \sqrt{\sigma u_\mu u^\mu} + \frac{e}{c} A_\mu u^\mu. \quad (16)$$

Hence, we see that the choice of sign for the relativistic Lagrangian (15) is intimately connected to the choice of space-time metric. We note that the components of the canonical four-momentum

$$p^\mu \equiv \begin{cases} \partial \mathcal{L}_\sigma / \partial \dot{x}_\mu = m \gamma \dot{x}^\mu + (e/c) A^\mu \\ \partial \mathcal{L}_\sigma / \partial u_\mu = m u^\mu + (e/c) A^\mu \end{cases} \quad (17)$$

are naturally independent of the choice ($\sigma = \pm 1$) of the space-time metric (1).

Lastly, the Lagrangian (15) satisfies the identity

$$\dot{x}^\mu \frac{\partial \mathcal{L}_\sigma}{\partial \dot{x}^\mu} = \left[m \gamma (\dot{x}^\mu \dot{x}_\mu) + \frac{e}{c} A_\mu \dot{x}^\mu \right] \equiv \mathcal{L}_\sigma, \quad (18)$$

where we used the definition (3) for the relativistic factor γ . Using the identity (18), the extended Hamiltonian $\mathcal{H}_\sigma$ derived from the extended Legendre transformation

$$\mathcal{H}_\sigma \equiv p_\mu \dot{x}^\mu - \mathcal{L}_\sigma \quad (19)$$

therefore vanishes identically.

In summary, while the choice for the sign ϵ of the relativistic Lagrangian (2) does not impact its associated Euler-Lagrange equations, the choice $\epsilon = -1$ is commonly used because of its natural connection with the relativistic Hamiltonian (4) and its non-relativistic limit (5). When a Lorentz-covariant relativistic Lagrangian formulation is adopted, however, a consistent choice for $\epsilon \equiv \sigma$ is associated with the sign σ of the space-time metric (1) used.

-
- [1] L. D. Landau and E. Lifshitz, *Classical Theory of Fields*, 4th ed. (Butterworth-Heinemann, 1980).
 - [2] J. D. Jackson, *Classical Electrodynamics*, 2nd ed. (Wiley, 1975).
 - [3] H. Goldstein, C. Poole, and J. Safko, *Classical Mechanics*, 3rd ed. (Addison Wesley, 2002).
 - [4] H. C. Corben and P. Stehle, *Classical Mechanics*, 2nd ed.

(Dover, 1960).

- [5] C. Lanczos, *The Variational Principles of Mechanics*, 4th ed. (Dover, 1970).
- [6] C. G. Gray, G. Karl, and V. A. Novikov, Rep. Prog. Phys. **67**, 159 (2004).